

RESEARCH ARTICLE

Open Access

# Characterization of Argentine honeys on the basis of their mineral content and some typical quality parameters

Marcelo Enrique Conti<sup>1\*</sup>, Maria Grazia Finoia<sup>2</sup>, Luca Fontana<sup>3</sup>, Giustino Mele<sup>1</sup>, Francesco Botrè<sup>4</sup> and Ivo Iavicoli<sup>3</sup>

## Abstract

**Background:** The levels of 19 elements (As, Be, Ca, Cd, Co, Cr, Cu, Fe, K, Mg, Mn, Na, Ni, Pb, Se, Tl, U, V, Zn) from sixteen different Argentine production sites of unifloral [eucalyptus (*Eucalyptus rostrata*), chilca (*Baccharis salicifolia*), Algarrobo (*Prosopis sp.*), mistol (*Ziziphus mistol*) and citric] and multifloral honeys were measured with the aim to test the quality of the selected samples. Typical quality parameters of honeys were also determined (pH, sugar content, moisture). Mineral elements were determined by using inductively coupled plasma mass spectrometer (ICP-MS DRC). We also evaluated the suitability of honey as a possible biomonitor of environmental pollution. Thus, the sites were classified through cluster analysis (CA) and then pattern recognition methods such as Principal Component Analysis (PCA) and discriminant analysis (DA) were applied.

**Results:** Mean values for quality parameters were: pH, 4.12 and 3.81; sugar 82.1 and 82.0 °brix; moisture, 16.90 and 17.00% for unifloral and multifloral honeys respectively. The water content showed good maturity. Likewise, the other parameters confirmed the good quality of the honeys analysed. Potassium was quantitatively the most abundant metal, accounting for 92,5% of the total metal contents with an average concentration of 832.0 and 816.2  $\mu\text{g g}^{-1}$  for unifloral and multifloral honeys respectively. Sodium was the second most abundant major metal in honeys with a mean value of 32.16 and 33.19  $\mu\text{g g}^{-1}$  for unifloral and multifloral honeys respectively. Mg, Ca, Fe, Mn, Zn and Cu were present at low-intermediate concentrations. For the other 11 trace elements determined in this study (As, Be, Cd, Co, Cr, Ni, Pb, Se, Tl, U and V), the mean concentrations were very low or below of the LODs. The sites were classified through CA by using elements' and physicochemical parameters data, then DA on the PCA factors was applied. Dendrograms identified three main groups. PCA explained 52.03% of the total variability with the first two factors.

**Conclusions:** In general, there are no evidences of pollution for the analysed honeys. The analytical results obtained for the Argentine honeys indicate the products' high quality. In fact, most of the toxic elements were below LODs. The chemometric analysis combining CA, DA and PCA showed their aptness as useful tools for honey's classification. Eventually, this study confirms that the use of honey as biomonitor of environmental contamination is not reliable for sites with low levels of contamination.

**Keywords:** Honey, Elements' content, Quality parameters, Multivariate statistical tools, Cluster analysis, Pattern recognition methods, Environmental biomonitoring

\* Correspondence: marcelo.conti@uniroma1.it

<sup>1</sup>Department of Management, University of Rome, Sapienza, Via del Castro Laurenziano 9, Rome 00161, Italy

Full list of author information is available at the end of the article

## Background

Honey is defined as “the natural sweet substance produced by *Apis mellifera* bees from the nectar of plants or from secretions of living parts of plants or excretions of plant-sucking insects on the living parts of plants, which the bees collect, transform by combining with specific substances of their own, deposit, dehydrate, store and leave in honeycombs to ripen and mature” [1]. Argentina is one of the major producers of bee honey and is the leading global exporter of high quality honey. About 50% come from the Province of Buenos Aires [2,3]. For instance, the United States imports 19% of their foreign honey from Argentina, spending about 54 million dollars in 2010. Also European countries such as Italy and Germany are strongly influenced by competition from Argentine and Chinese varieties whose prices are lower by roughly 50% [4].

Honey has high nutritional value (330 kcal/100 g) and fast absorption of its carbohydrates on consumption. It is a high carbohydrate food and shows anti-bacterial and anti-inflammatory properties in the treatment of skin wounds and several gastrointestinal diseases [4-10]. Honey activates the immune system and its ingestion may be beneficial with respect to cancer and metastasis prevention [5,10]. Hydrogen peroxide produced enzymatically is responsible for honey's antibacterial activity [7]. Honey is a potent inhibitor of the bacterium *Helicobacter pylori* that causes peptic ulcers and gastritis [10].

The European Union [1] defines general and specific compositional characteristics of honey such as sugar content, humidity, acidity, electrical conductivity, diastase activity and hydroxymethylfurfural (HMF) content. Moreover, labels on honey packaging should report information on the product's regional or topographical origin, floral or vegetable origin. If honey originates from different countries the label should specify “blend of EU (or no-EU) honeys” or “blend of EU and no-EU honeys”.

The composition and properties of a particular honey sample depend highly on the type of flowers visited by the bees, as well as on the climatic conditions in which the plants grow [11,12]. Melissopalynology (pollen analysis) is the traditional method used to determine the botanical origin of honeys [13,14], but this technique has some limitations [15]. In fact, melissopalynology requires relevant knowledge of pollen morphology and specialised professional personnel to achieve reliable results [13]. However, nowadays in spite of these problems melissopalynology remains the reference method.

Bees forage an area of about 7 km<sup>2</sup> and come in contact constantly with the surrounding environment. The chemical composition and properties of honey depend on the type of flowers visited by the bees, as well as on the climatic conditions in which the plants grow [4,9]. This also implies that honeybees and their products (i.e.

pollen, wax, etc.) can be employed as potential biomonitoring of environmental contamination [9,16]. Finally, the specific chemical and physical properties can be used for the determination of the botanical origin of honey [17,18].

The role of elements in honey is of high relevance [19-21] in terms of both its quality and safety. Scarce information is available on the elements' composition of Argentine honeys, this is also connected with the relevance of Argentina as one of the main honey exporter in the world. In fact, only 5% of the total honey production in Argentina is destined to domestic consumption.

The aim of the work was to measure the levels of 19 elements (As, Be, Ca, Cd, Co, Cr, Cu, Fe, K, Mg, Mn, Na, Ni, Pb, Se, Tl, U, V, Zn) and some typical quality parameters (pH, sugar content, moisture) from sixteen different Argentine production sites of unifloral [i.e. eucalyptus, chilca (*Baccharis* spp.), Algarrobo (*Prosopis* sp.), mistol (*Ziziphus mistol*) and citric] and multifloral honey samples. Duplicate samples were taken from each production area. Due to its economic relevance for export purposes and for production levels, samples were mainly collected in the Province of Buenos Aires.

We have evaluated, by means of multivariate statistical methods whether the physicochemical parameters and the elements' content can classify or discriminate the sampling sites in order to confirm the suitability of honey as a possible biomonitor of environmental pollution. Thus, the sites were classified through cluster analysis by using elements' data, then pattern recognition methods such as Principal Component Analysis (PCA) and discriminant analysis (DA) were applied (see Statistical methods section for details).

## Results and discussion

Table 1 reports the mean  $\pm$  SD and the range of the physicochemical parameters for the analysed honeys. Reported data for the physicochemical parameters were homogeneous for the analyzed honeys, showing very low SD levels (Table 1).

**Table 1 Descriptive statistics for physicochemical parameters in Argentine honey samples**

	N	Mean $\pm$ SD	MIN	MAX
<b>SUGAR [°brix]</b>				
Unifloral	5	82.1 $\pm$ 0.7	81.5	83.0
Multifloral	11	82.0 $\pm$ 1.2	80.5	84.0
<b>pH</b>				
Unifloral	5	4.12 $\pm$ 0.21	3.87	4.46
Multifloral	11	3.81 $\pm$ 0.27	3.55	4.43
<b>MOISTURE [%]</b>				
Unifloral	5	16.9 $\pm$ 0.6	16.0	17.5
Multifloral	11	17.0 $\pm$ 1.3	15.0	19.0

The pH ranged between 3.87 and 4.46 with a mean value of 4.12 for unifloral honeys, while the range was between 3.55 and 4.43 with a mean of 3.81 for multifloral honeys.

Most bacteria and moulds grow in a neutral and mildly alkaline environment respectively, while yeasts require an acidic environment (pH = 4.0 – 4.5) and do not grow in alkaline media [22]. In fact, pH is a useful index of possible microbial contamination [23] and has great relevance during the extraction and storage of honey because it is connected with the texture and the product's shelf life [24].

The mean pH value (3.81) of multifloral Argentine honeys was similar than those reported by Baroni et al. [25] for Córdoba (Argentina) honeys and by Conti et al. [4] for Italian honeys (Marche Region); while the obtained mean pH value for unifloral honeys (i.e. 4.12) was comparable to those of Lazio Region (Italy) honeys [24] and to Andalusian (Spain) unifloral honeys [26].

Water content is connected with the climatic conditions and the degree of maturity; anomalous values may be an index of adulterations. It generally depends on the botanical origin of the sample, the processing techniques and the storage conditions [4,24,27]. Mean humidity was 17.0% with a range of between 15.0 – 19.0% for multifloral honeys and a mean of 16.9% with a range of between 16.0 – 17.5% for unifloral honeys. None of the samples exceeded the limit permitted of 20% by the Codex Alimentarius [28] and the Council Directive [1]. However one sample (code M8, Entre Rios Province) showed a value higher than the limit of 18% established by the Argentine legislation [29]. Our results generally confirm that the fermentation rate is very low in the analyzed samples. Moisture values observed for our samples were slightly lower than those obtained for Córdoba (Argentina) honeys [25] and quite similar to honeys from Buenos Aires Region [30], and to those of Andalusia (Spain) [26], Lazio Region (Italy) [24], Serbian Acacia honeys [27] and to *Entre-Douro e Minho* Region of Portugal [31].

The average total sugar content was 82.1% and the range was 81.5 – 83.0% for unifloral honeys, and a mean of 82.0% with a range of between 80.5 – 84.0% for multifloral honeys. Our results match those obtained for Italian and Spanish honeys [4,23,24,26].

Tables 2 and 3 show the mean, standard deviation and the concentration ranges of the elements for unifloral and multifloral honey samples respectively.

Potassium was quantitatively the most abundant metal, accounting for 92,5% of the total metal contents with an average concentration of 832.0 and 816.2  $\mu\text{g g}^{-1}$  (wet weight) for unifloral and multifloral Argentine honeys respectively. For data comparison, the results were appropriately transformed (i.e. wet or dry basis) when necessary.

As above pointed out, little information is available on mineral content in Argentine honeys. Our mean potassium levels were considerably higher than those of Córdoba (Argentina) honeys [25] and to Lazio (Italy) honeys [24] and higher to Marche region honeys (Italy) [4]. This result is consistent with other reported data [9,19,32].

Sodium, as expected, was the second most abundant major metal in honeys. We determined a mean sodium content of 32.16 and 33.19  $\mu\text{g g}^{-1}$  (w.w.) for unifloral and multifloral Argentine honeys respectively. Mean sodium values were lower than Córdoba (Argentina) honeys [25] and Lazio (Italy) honeys [24]; while we obtained slightly higher mean values than those reported for Marche region (Italy) honeys [4].

Mean magnesium levels were 17.26 and 22.64  $\mu\text{g g}^{-1}$  w.w. in unifloral and multifloral Argentine honeys respectively, and were quite higher than those determined in Córdoba (Argentina) honeys [25]. Calcium mean levels, i.e. 6.92 and 10.86  $\mu\text{g g}^{-1}$  w.w. for unifloral and multifloral Argentine honeys respectively, were lower than Córdoba (Argentina) honeys [25], than to Spanish [33], and to Lazio and Marche Regions (Italy) honeys [4,24]. The mean iron levels in our samples, 3.57 and 2.99  $\mu\text{g g}^{-1}$  w.w. for unifloral and multifloral Argentine honeys respectively, were lower than those of Córdoba (Argentina) [25], to San Luis – La Pampa honeys (middle Argentina) [34] and to Lazio and Marche Regions (Italy) honeys [4,24].

The mean manganese levels (2.61 and 1.05  $\mu\text{g g}^{-1}$  w.w.) were higher to those found for Córdoba (Argentina) honeys [25], and to Marche region (Italy) honeys [4], and lower than Lazio region (Italy) honeys [24]. The mean zinc levels (0.87 and 1.17  $\mu\text{g g}^{-1}$  w.w.) were lower than those found for Córdoba (Argentina) honeys [25] and Lazio honeys [24] and comparable to those found for San Luis – La Pampa honeys (middle Argentina) [34]. The mean copper levels for our samples (0.18 and 0.29  $\mu\text{g g}^{-1}$ ) are lower than those of Lazio honeys [24] while Cu was not detected in the Córdoba (Argentina) honeys [25].

Potassium showed positive correlation with Ca [ $r = 0.70$  t(14) = 3.68 p = 0.002], Mg [ $r = 0.76$  t(14) = 4.44 p < 0.001] and Cu [ $r = 0.73$  t(14) = 4.04 p = 0.001]. Calcium correlated positively with Mg [ $r = 0.77$  t(14) = 4.53 p < 0.001] and Cu [ $r = 0.50$  t(14) = 2.17 p = 0.047]. Sodium correlated positively with Ca [ $r = 0.524$  t(14) = 2.30 p = 0.037].

The presence of these major essential elements in honey such as Ca, K, Na and Mg is of certain nutritional relevance and dietary value, mainly connected with children's health [9,19]. As previously reported [24] the intake of many major and minor metals (i.e. Fe, Mn, Cu and Zn) from honey is very low due to its low level of consumption. Generally, honey does not contribute for a significant proportion of minerals recommended dietary allowances (RDAs), usually from a few per cent, or even lower [24,35].

**Table 2 Descriptive statistics of elements' content ( $\mu\text{g g}^{-1}$  wet weight) in Argentine unifloral honey samples**

	Eucalipto (EntreRios)	Chilca (Córdoba)	Algarrobo (S. del Estero)	Mistol (S. del Estero)	Citric (Tucumán)	Mean $\pm$ SD	Min	Max
	M1	M2	M3	M4	M5			
<b>As</b>	<0.01	<0.01	<0.01	<0.01	<0.01	-	<0.01	<0.01
<b>Be</b>	<0.01	<0.01	<0.01	<0.01	<0.01	-	<0.01	<0.01
<b>Ca</b>	6.18	2.50	4.13	13.74	8.04	6.92 $\pm$ 4.35	2.50	13.74
<b>Cd</b>	<0.01	<0.01	<0.01	0.01	<0.01	0.01 $\pm$ 0.00	<0.01	0.01
<b>Co</b>	0.01	<0.01	<0.01	<0.01	<0.01	0.01 $\pm$ 0.00	<0.01	0.01
<b>Cr</b>	<0.01	<0.01	<0.01	<0.01	<0.01	-	<0.01	<0.01
<b>Cu</b>	0.27	0.12	0.20	0.18	0.15	0.18 $\pm$ 0.06	0.12	0.27
<b>Fe</b>	3.38	2.19	4.05	3.73	4.50	3.57 $\pm$ 0.87	2.19	4.50
<b>K</b>	488.4	251.1	318.4	2022.6	1079.8	832.0 $\pm$ 741.3	251.1	2022.6
<b>Mg</b>	14.37	4.59	8.12	37.98	21.26	17.26 $\pm$ 13.21	4.59	37.98
<b>Mn</b>	8.84	0.77	1.14	1.60	0.70	2.61 $\pm$ 3.50	0.70	8.84
<b>Na</b>	62.00	7.21	39.27	34.65	17.67	32.16 $\pm$ 21.09	7.21	62.00
<b>Ni</b>	0.05	0.01	0.06	0.02	0.03	0.04 $\pm$ 0.02	0.01	0.06
<b>Pb</b>	0.01	0.02	0.02	0.01	0.02	0.02 $\pm$ 0.01	0.01	0.02
<b>Se</b>	0.01	<0.01	<0.01	0.01	<0.01	0.01 $\pm$ 0.00	0.01	0.01
<b>Tl</b>	<0.01	<0.01	<0.01	<0.01	<0.01	-	<0.01	<0.01
<b>U</b>	<0.01	<0.01	<0.01	<0.01	<0.01	-	<0.01	<0.01
<b>V</b>	<0.01	<0.01	<0.01	<0.01	<0.01	-	<0.01	<0.01
<b>Zn</b>	0.55	1.05	1.42	0.84	0.51	0.87 $\pm$ 0.38	0.51	1.42

For the other 11 trace elements determined in this study (As, Be, Cd, Co, Cr, Ni, Pb, Se, Tl, U and V), the mean concentrations were very low or below of the LODs (Tables 2 and 3). In general, the presence of these elements can indicate contamination during honey processing, shipping or storage connected with the use of steel or galvanized containers [19]. However, in an another study chromium levels in some honey samples collected in the Buenos Aires province (Argentina) were in the range of 0.9 – 6  $\mu\text{g g}^{-1}$  [36] while a mean concentration of 0.47  $\mu\text{g g}^{-1}$  is reported for San Luis – La Pampa honeys (middle Argentina) [34].

Lead levels were lower than others obtained for honeys collected in the Buenos Aires province (Argentina) [36], and comparable to those reported for Turkish honeys [37], while our nickel levels were higher than those of Turkish honeys [37]. Generally, with rare exceptions, our data for some trace elements (i.e. As, Cd, Co, Cr, Pb) are comparable or at a lower levels than those reported for other countries [38-42]. Very scarce information is present in literature about some elements in Argentine honeys such as Ni, Se, Tl, U and V. Our selenium levels were lower than those measured in Turkish honeys [37]. While we not detected uranium and vanadium in our samples (LOD < 0.01  $\mu\text{g g}^{-1}$ , Tables 2 and 3) Almeida-Silva et al. [43] reported levels of 0.34 and 0.28  $\mu\text{g g}^{-1}$  of U and 13.5 and 5.6  $\mu\text{g g}^{-1}$  of vanadium in Portuguese honeys.

We have no found data for beryllium in honeys in the literature. However, we detected Be only in the Entre Ríos (M8) multifloral sample (0.10  $\mu\text{g g}^{-1}$ ) while the others were below LODs. At present, there are no studies that suggested a health risk for the presence of beryllium in food and drinking water, even if it can be an index of industrial contamination [44].


Overall, from our results we can infer that there are no evidences of pollution for the analysed honeys and these results confirm their good quality.

We applied the hierarchical clustering by using data set (Tables 1, 2 and 3, see also Experimental section below) in order to classify the production sites. The result of the cluster analysis is reported in Figure 1. The x axis depicts the sampling sites while the y axis indicates the calculated distances among sampling sites. From Figure 1 we observe that there are three main groups. The honeys produced in the provinces of Córdoba and Tucumán are grouped in the A cluster. The honeys produced in the provinces of Entre Ríos and Santiago (i.e. Santiago del Estero) are grouped in the B cluster, and those produced in the Buenos Aires province are linked with the A cluster at the last step of the iterative aggregation process.

Thus, in order to discriminate the elements and parameters classified by cluster analysis, we applied discriminant analysis (DA) on principal component analysis (PCA) factors (see experimental section). PCA explained 52.03%

**Table 3 Descriptive statistics of elements' content ( $\mu\text{g g}^{-1}$  wet weight) in Argentine multifloral honey samples**

	Córdoba	Prov. Buenos Aires	Entre Ríos	Entre Ríos	Prov. Buenos Aires	Mean $\pm$ SD	Min	Max						
	M6	M7	M8	M9	M10	M11	M12	M13	M14	M15	M16			
<b>As</b>	<0.01	<0.01	<0.01	0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.01 $\pm$ 0.00	<0.01	0.01
<b>Be</b>	<0.01	<0.01	0.10	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.10 $\pm$ 0.00	<0.01	0.10
<b>Ca</b>	7.57	6.03	17.26	18.97	15.33	14.68	13.61	18.61	2.93	1.97	2.50	10.86 $\pm$ 6.75	1.97	18.97
<b>Cd</b>	<0.01	<0.01	<0.01	0.03	0.01	0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.02 $\pm$ 0.01	<0.01	0.03
<b>Co</b>	<0.01	<0.01	0.01	0.01	<0.01	0.01	<0.01	<0.01	<0.01	0.01	<0.01	0.01 $\pm$ 0.00	<0.01	0.01
<b>Cr</b>	<0.01	0.01	0.01	<0.01	<0.01	<0.01	0.05	0.04	<0.01	<0.01	<0.01	0.03 $\pm$ 0.02	<0.01	0.05
<b>Cu</b>	0.13	0.09	0.38	1.19	0.17	0.20	0.12	0.32	0.20	0.17	0.21	0.29 $\pm$ 0.31	0.09	1.19
<b>Fe</b>	4.40	2.07	3.15	2.24	2.32	4.18	2.26	3.99	2.40	3.22	2.68	2.99 $\pm$ 0.86	2.07	4.40
<b>K</b>	1225.8	324.9	1408.2	2813.3	755.8	476.2	412.3	1107.5	176.6	134.1	143.9	816.2 $\pm$ 802.6	134.1	2813.3
<b>Mg</b>	18.12	12.39	75.38	49.77	19.68	23.57	16.46	23.87	3.43	3.01	3.31	22.64 $\pm$ 21.95	3.01	75.38
<b>Mn</b>	0.57	0.55	2.39	3.13	1.18	0.57	0.14	0.40	0.60	1.40	0.59	1.05 $\pm$ 0.93	0.14	3.13
<b>Na</b>	20.05	4.88	36.30	28.62	105.95	25.19	25.79	76.44	13.26	15.45	13.16	33.19 $\pm$ 30.66	4.88	105.95
<b>Ni</b>	0.02	0.02	0.07	0.05	0.02	0.05	0.03	0.04	0.03	0.05	0.02	0.04 $\pm$ 0.02	0.02	0.07
<b>Pb</b>	0.01	0.04	0.01	0.01	0.04	0.04	0.01	0.04	0.01	0.03	0.01	0.02 $\pm$ 0.02	0.01	0.04
<b>Se</b>	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	-	<0.01	<0.01
<b>Tl</b>	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	-	<0.01	<0.01
<b>U</b>	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	-	<0.01	<0.01
<b>V</b>	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	-	<0.01	<0.01
<b>Zn</b>	2.75	1.22	0.80	0.70	0.81	2.04	1.50	0.74	0.72	0.92	0.70	1.17 $\pm$ 0.67	0.70	2.75



of the total variability with the first two factors. Elements such as As, Be, Cd, Co, Cr, Se, Tl, U, V had to be eliminated from the data set because of their too low levels (or below LODs). Therefore, the data matrix was constituted by thirteen loadings: Ca, Cu, Fe, K, Mg, Mn, Na, Ni, Pb, Zn, moisture, sugar content and pH. Results are reported in Figure 2.

From this study we can draw some findings. First, the cluster A (honeys from Córdoba and Tucumán provinces) has higher concentrations of Zn and Fe and lower concentrations of Ca, Na and Ni compared to the other groups. Second, the cluster B (Entre Ríos and Santiago honeys) is characterized by higher concentrations of K, Mg, Mn, Cu and Ni and higher pH levels than the other clusters. Moreover, the honeys of the cluster B have lower levels of Zn than the others. Third, the cluster C (Buenos Aires province honeys) showed higher levels of Pb than the other groups and lower levels of sugar, pH and K, Mg, Mn and Fe.

However, these results can be considered as indicative because the DA did not result significant (Montecarlo test  $RV = 0.126$   $p = 0.754$ ). Thus, the elements and physiological parameters determined in honey have weak discriminating power. This is of relevance because it further confirms our previous statements [16] that honey is not a reliable biomonitor of environmental contamination, in particular in sites with low levels of contamination (see also refs. [9,45] for discussion). Furthermore, we also conducted the DA on the PCA factors considering the variables 'unifloral' and 'multifloral' honeys for the elements and physiological parameters determined (results not shown), but also in this case DA resulted not significant (Montecarlo test  $RV = 0.073$   $p = 0.309$ ). This further agrees with the statements above reported.

## Conclusions

In general, we can infer that there are no evidences of pollution for the analysed honeys. The analytical results obtained for the Argentine honeys indicate the products' high quality. In fact, most of the toxic elements were below LODs. The chemometric analysis combining CA, DA and PCA showed their aptness as useful tools for honey's classification. This study further confirms that the use of honey as biomonitor of environmental contamination is not reliable for sites with low levels of contamination.

Eventually, further research is needed in order to characterize Argentine honeys by means of pattern recognition methods of zones with high production levels and in order to improve their economic interest.

## Experimental

### Samples


The study was conducted on 16 samples of the typical honeys coming from different production areas in Argentina. Five unifloral and eleven multifloral samples were collected and analysed in duplicate (see Table 4 for description). All collected samples were taken from the local beekeepers' association with a guarantee of genuineness. All samples were collected, stored in plastic holders and kept at 4–5°C until analysis.

### pH, sugar content and moisture

The pH was assessed by means of a potentiometer utilizing a pH meter Mettler Delta 345 (Mettler Toledo, Milano, Italy) [46]. Sugar and moisture values were determined utilizing a Bertuzzi refractometer (Bertuzzi, Milano, Italy) owing two direct reading displays, for the measurement of sugar content and moisture percent respectively (Chatway method). Total sugar content was expressed as brix degrees [46].

### Determination of mineral elements

About 0.8 g of fresh honey was treated with 3.5 ml of 70% (w/w) Nitric Acid Suprapur (Merck, Suprapur, Darmstadt, Germany) and 1.5 ml of 30% (w/w) Hydrogen Peroxide Suprapur (Merck, Darmstadt, Germany) in PTFE vessels. The microwave closed digestion system (Milestone, Start D) was used for the mineralization process. The treatment procedure was programmed in four steps with a power of 1200 W applied for 5, 3, 8 and 15 min each respectively. The temperature was 120°C for the first two steps and 200°C for the second two steps. A sample of reference materials and blank was included in each analytical batch. Subsequently, digestion vessels were cooled to room temperature. The final clear solution was made up to 15 mL with DWI water. Digestion methods in biological and environmental matrices were discussed in our previous studies [47,48].



The samples were quantified for 19 elements: As, Be, Ca, Cd, Co, Cr, Cu, Fe, K, Mg, Mn, Na, Ni, Pb, Se, Tl, U, V and Zn, in digested honeys by inductively coupled plasma mass spectrometry [ICP-MSDRC-e (Dynamic Reaction Cell) mod. Elan, Perkin Elmer]. Traceability of results was obtained from the analysis of the certified reference materials NIST-1515 (Apple leaves - National Institute of Standards and Technology) and the Antarctic Krill MURST-ISS-A2 (Italian Research Programme in Antarctica). For the MURST-ISS-A2 the mean recovery percentages (five replicates) were: As:  $94.1 \pm 2.8\%$ ; Cd:  $93.6 \pm 2.8\%$ ; Co:  $101.2 \pm 1.8\%$ ; Cr:  $98.0 \pm 1.1\%$  (not certified, informative concentration); Cu:  $101.4 \pm 2.0\%$ ; Ni:  $97.9 \pm 2.4\%$ ; Pb:  $97.0 \pm 0.9\%$ ; Se:  $96.8 \pm 2.9\%$  and Zn:  $102.1 \pm 2.8\%$ . For the NIST-1515 the mean recovery percentages (five replicates) were: Ca:  $101.5 \pm 2.9\%$ ; Fe  $100.9 \pm 2.9\%$ ; K:  $98 \pm 1.8\%$ ; Mg:  $102.2 \pm 2.7\%$ ; Mn:  $98 \pm 2.2\%$ ; Na  $99.5 \pm 1.8\%$  and V:  $101.2 \pm 2.8\%$ . Results were in very good agreement with certified values for the tested elements proving good accuracy of the method employed.

All chemicals used in sample treatment were ultra-pure grade ( $\text{HNO}_3$ ,  $\text{H}_2\text{O}_2$  30%, Merck, Suprapur, Darmstadt, Germany). Ultra-pure water (Milli-Q system, Millipore Corporation, U.S.A.) was used for all solutions. All glass-ware was cleaned prior to use by soaking in 10% v/v  $\text{HNO}_3$  for 24 hours before rinsing with Milli-Q water. The standard metal solutions were prepared from stock standard solutions of ultra-pure grade supplied by Merck (Darmstadt, Germany).

The laboratory precision for the whole analytical process was tested by measuring the Relative Standard Deviation (RSD %) of ten replicates for each sample tested ( $n = 10$ ). The obtained values for the analysed elements were always below 7% proving the good repeatability of the analytical method. For details about uncertainty, precision, accuracy and methods validation see refs [47-50].

#### Statistical methods

Duplicate honey samples were collected at each site. The normality of distributions of the 19 elements determined

**Table 4 Argentine honey samples description**

Samples	Product	Geographical origin
M1	Eucalipto honey	Colón, Entre Ríos
M2	Chilca honey	Sierras de Córdoba, Córdoba
M3	Algarrobo honey	Aguirre, Santiago del Estero
M4	Mistol honey	Aguirre, Santiago del Estero
M5	Citric honey	Alberdi, Tucumán
M6	Multifloral honey (organic)	Cruz del eje, Córdoba
M7	Multifloral honey (pasture)	Castelar, Province of Buenos Aires
M8	Multifloral honey	Guauguaychú, Entre Ríos
M9	Multifloral honey	Concordia, Entre Ríos
M10	Multifloral honey	Mar del Plata, Province of Buenos Aires
M11	Multifloral honey	Berazategui, Province of Buenos Aires
M12	Multifloral honey (creamed)	La Plata, Province of Buenos Aires
M13	Multifloral honey (organic)	Central area, Province of Buenos Aires
M14	Multifloral honey	Tandil, Province of Buenos Aires
M15	Multifloral honey	Central area, Province of Buenos Aires
M16	Multifloral honey	Sierra de los Padres, Mar del Plata, Province of Buenos Aires

was tested by using Shapiro-Wilk test [51,52]. Then the homogeneity for the data of the dual collected samples was tested. The t Student test was applied for testing univariate paired comparisons when normal distribution was obtained and the non parametric Wilcoxon rank sum test was applied for the other elements with not normal distribution [53,54]. Results confirmed the data homogeneity (data not shown) and then we merge the data of duplicated samples.

Several approaches can be employed for data analysis in this kind of studies [55,56]. In this work we first applied the hierarchical cluster analysis [57-60] to the data set (16 samples; and 19 elements plus 3 physiological parameters) in order to classify the distribution of the honey samples according to their production areas. The optimal number of clusters was determined by using the hierarchical cut-clustering rule.

The hierarchical clustering by minimum (energy) E-distance method was performed. Dissimilarities are  $\|x-y\|^a$  where the exponent is in the interval (0,2). This function performs agglomerative hierarchical clustering. Initially, each of the n singletons is a cluster. At each of n-1 steps, the procedure merges the pair of cluster with minimum E-distance. The E-distance between two cluster  $C_i$ ,  $C_j$  of size  $n_i$  and  $n_j$  is given by

$$e(C_i, C_j) = \frac{n_i n_j}{n_i + n_j} [2M_{ij} - M_{ii} - M_{jj}]$$

where

$$M_{ij} = \frac{1}{n_i n_j} \sum_{p:1}^{n_i} \sum_{q:1}^{n_j} \|X_{ip} - X_{jq}\|^a$$

$\| \cdot \|$  denotes Euclidean norm,  $X_{ip}$  denotes the p-th observation in the i-th cluster.

Then, discriminant analysis (DA) on principal component analysis (PCA) factors [61] was conducted with the aim to discriminate the elements and parameters classified by cluster analysis (CA). The 13 factors extracted by PCA (i.e. 100% of information) were included in DA. The analysis' significance was tested by Monte Carlo test (non parametric version of the Pillai's test) based on 999 replicates [62].

DA and PCA are considered statistical tools capable to reveal structures in environmental data [55,63,64]. In particular, PCA is a statistic technique belonging to the so-called "unsupervised pattern recognition methods," useful for carrying out exploratory data analysis when there is no preliminary knowledge about the characteristics (i.e., distribution and structure) of the data to be analyzed [65]. DA is a statistic technique belonging to the so-called "supervised pattern recognition methods," useful for carrying out specific data analysis when a previous unsupervised pattern recognition method, such as PCA, has suggested a potential discrimination among the data. Mainly for this cause, DA is applied to PCA results [66].

Data analysis was performed using the software R version 2.15.2 (2012-10-26) - "Trick or Treat" and the packages Energy, Ade4.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

MEC conceived of the study and, together with FB, MGF and II participated in its design and drafted the manuscript. MEC, GM and LF coordinated the sampling protocols and the whole analytical procedures. MEC and MGF participated in the design and performed the statistical analysis. This project was based on the ideas and under the guidance and consultation of MEC, MGF, FB and II. All authors read and approved the final manuscript.

### Acknowledgements

M.E. Conti wishes to thank Cecilia Gauna for her assistance in the sample collection and classification.

### Author details

<sup>1</sup>Department of Management, University of Rome, Sapienza, Via del Castro Laurenziano 9, Rome 00161, Italy. <sup>2</sup>Italian National Institute for Environmental Protection and Research, Viale V. Brancati 60, Rome 00166, Italy. <sup>3</sup>Institute of Occupational Medicine, Università Cattolica del Sacro Cuore, Largo Francesco Vito 1, Rome 00168, Italy. <sup>4</sup>Department of Experimental Medicine, University of Rome, Sapienza, Viale Regina Elena 324, Rome 00161, Italy.

Received: 3 April 2014 Accepted: 24 June 2014

Published: 2 July 2014

### References

1. Council Directive of the European Union: Council Directive 2001/110/EC of 20 December 2001 relating to honey. *Offic J Eur Communities* 2002, **L10**:47–52.
2. Domínguez MA, Grünhut M, Pistonesi MF, Di Nezio MS, Centurión ME: Automatic flow-batch system for cold vapor atomic absorption spectroscopy determination of mercury in honey from Argentina using online sample treatment. *J Agric Food Chem* 2012, **60**:4812–4817.
3. Finola-Mónica S, Lasagno-Mirta C, Marioli-Juan M: Microbiological and chemical characterization of honeys from central Argentina. *Food Chem* 2007, **100**(4):1649–1653.
4. Conti ME, Stripeikis J, Campanella L, Cucina D, Tudino MB: Characterization of Italian honeys (Marche Region) on the basis of their mineral content and some typical quality parameters. *Chem Cent J* 2007, **1**:14.
5. Bogdanov S, Jurendic T, Sieber R, Gallmann P: Honey for nutrition and health: a review. *Am J Coll Nutr* 2008, **27**:677–689.
6. Greenwood D: Honey for superficial wounds and ulcers. *Lancet* 1993, **341**(8837):90–91.
7. Taormina PJ, Niemira BA, Beuchat LR: Inhibitory activity of honey against food borne pathogens as influenced by the presence of hydrogen peroxide and level of antioxidant power. *Int J Food Microbiol* 2001, **69**:217–225.
8. Molan P: Why honey is effective as a medicine: 2. The scientific explanation of its effects. *Bee World* 2001, **82**:22–40.
9. Pohl P: Determination of metal content in honey by atomic absorption and emission spectrometries. *Trend Anal Chem* 2009, **28**(1):117–128.
10. Alvarez-Suarez JM, Tulipani S, Romandini S, Bertoli E, Battino M: Contribution of honey in nutrition and human health: a review. *J Nutr Metab* 2010, **3**:15–23.
11. Abu-Tarboush HM, Al-Kahtani HA, El-Sarrage MS: Floral-type identification and quality evaluation of some honey types. *Food Chem* 1993, **46**:13–17.
12. Perez-Arquillue C, Conchello P, Arino A, Juan T, Herresa A: Quality evaluation of Spanish rosemary (*Rosmarinus officinalis*) honey. *Food Chem* 1994, **51**:207–210.
13. Louveaux J, Maurizio A, Vorwohl G: Methods of melissopolynology. *Bee World* 1978, **59**:139–157.
14. Song X-Y, Yao Y-F, Yang W-D: Pollen Analysis of Natural Honeys from the Central Region of Shanxi, North China. *PLoS Biol* 2012, **7**(11):e49545.
15. Molan PC: The limitations of the methods of identifying the floral source of honeys. *Bee World* 1998, **79**:59–68.
16. Conti ME, Botrè F: Honeybees and their products as potential bioindicators of heavy metals contamination. *Environ Mon Ass* 2001, **69**:267–282.
17. Latorre MJ, Peña R, Pita C, Botana A, García S, Herrero C: Chemometric classification of honeys according to their type. II. Metal content data. *Food Chem* 1999, **66**:263–268.
18. Bogdanov S, Ruoff K, Persano Oddo L: Physico-chemical methods for the characterisation of unifloral honeys: a review. *Apidologie* 2004, **35**:S4–S17.
19. Pohl P, Stecka H, Sergiel I, Jamroz P: Different Aspects of the Elemental Analysis of Honey by Flame Atomic Absorption and Emission Spectrometry: A Review. *Food Anal Methods* 2012, **5**:737–751.
20. Bogdanov S, Haldimann M, Luginbuhl W, Gallmann P: Minerals in honey: environmental, geographical and botanical aspects. *J Apic Res Bee World* 2007, **46**:269–275.
21. Pohl P, Sergiel I, Stecka H: Determination and fractionation of metals in honey. *Crit Rev Anal Chem* 2009, **39**:276–288.
22. Conti ME, Saccares S, Cubadda F, Cavallina R, Tenoglio CA, Ciprotti L: Il miele nel Lazio: indagine sul contenuto in metalli in tracce e radionuclidi. *Riv Sci Alim* 1998, **2**:107–119.
23. Terrab A, Recamales AF, Hernanz D, Heredia FJ: Characterisation of Spanish thyme honeys by their physicochemical characteristics and mineral contents. *Food Chem* 2004, **88**:537–542.
24. Conti ME: Lazio region (central Italy) honeys: determination of mineral content and typical quality parameters. *Food Contr* 2000, **11**:459–463.
25. Baroni MV, Arrua C, Nores ML, Fayé P, Diaz MDP, Chiabrando GA, Wunderlin DA: Composition of honey from Córdoba (Argentina): Assessment of North/South provenance by chemometrics. *Food Chem* 2009, **114**:727–733.
26. Serrano S, Villarejo M, Espejo R, Jodral M: Chemical and physical parameters of Andalusian honey: Classification of Citrus and Eucalyptus honeys by discriminant analysis. *Food Chem* 2004, **87**:619–625.
27. Lazarevic KB, Andric F, Trifkovic J, Tesik Z, Milojkovic-Opsenica D: Characterization of Serbian unifloral honeys according to their physicochemical parameters. *Food Chem* 2012, **132**:2060–2064.
28. Codex Alimentarius Commission: Revised Codex Standard for Honey Codex Stan 12–1981, Rev. 1 (1987), Rev. 2 (2001). *Codex Standard* 1981, **12**:1–7.
29. Código Alimentario Argentino (CAA): *Productos azucarados*, Volume 10. De La Canal y Asociados S.R.L.; 1997:782–785.
30. Silvano MF, Varela MS, Palacio MA, Ruffinengo S, Yamul DK: Physicochemical parameters and sensory properties of honeys from Buenos Aires region. *Food Chem* 2014, **152**:500–507.
31. Feas X, Pires J, Estevinho ML, Iglesias A, Araujo JPP: Palynological and physicochemical data characterisation of honeys produced in the Entre-Douro e Minho region of Portugal. *Int J Food Sci Tech* 2010, **45**:1255–1262.
32. Silva LR, Videira R, Monteiro AP, Valentao P, Andrade PB: Honey from Luso region (Portugal). Physicochemical characteristics and mineral content. *Talanta* 2009, **93**:73–77.
33. Fernández-Torres R, Pérez-Bernal JL, Bello-López MÁ, Callejón-Mochón M, Jiménez-Sánchez JC, Guiraúm-Pérez A: Mineral content and botanical origin of Spanish honeys. *Talanta* 2005, **65**:686–691.
34. Pellerano RG, Unates MA, Cantarelli MA, Camina JM, Marchevsky EJ: Analysis of trace elements in multiflora Argentine honeys and their classification according to provenance. *Food Chem* 2012, **134**:578–582.
35. Grembecka M, Szefer P: Evaluation of honeys and bee products quality based on their mineral composition using multivariate techniques. *Environ Monit Assess* 2013, **185**:4033–4047.
36. Enrich C, Boeykens S, Caracciolo N, Custo G, Vázquez C: Honey characterization by total reflection X-ray fluorescence: evaluation of environmental quality and risk for the human health. *X-Ray Spectrom* 2007, **36**:215–220.
37. Tuzen M, Silici S, Mendil D, Soyulak M: Trace element levels in honeys from different regions of Turkey. *Food Chem* 2007, **103**:325–330.
38. Chudzinska M, Baralkiewicz D: Estimation of honey authenticity by multielements characteristics using inductively coupled plasma-mass spectrometry (ICP-MS) combined with chemometrics. *Food Chem Toxicol* 2010, **48**:284–290.
39. Vanhanen LP, Emmertz AB, Savage GP: Mineral analysis of mono-floral New Zealand honey. *Food Chem* 2011, **128**:236–240.
40. Habib HM, Al Meqbali FT, Kamal H, Souka UD, Ibrahim WH: Physicochemical and biochemical properties of honeys from arid regions. *Food Chem* 2014, **153**:35–43.
41. Bilandzic N, Dokic' M, Sedak M, Solomun Kolanovic' B, Varenina I, Koncurat A, Rudan N: Determination of trace elements in Croatian floral honey originating from different regions. *Food Chem* 2011, **128**:1160–1164.
42. Bilandzic N, Gacic M, Dokic M, Sedak M, Sipusic DI, Koncurat A, Gajger IT: Major and trace elements levels in multifloral and unifloral honeys in Croatia. *J Food Comp Anal* 2014, **33**:132–138.
43. Almeida-Silva M, Canha N, Galinha C, Dung HM, Freitas MC, Siteo T: Trace elements in wild and orchard honeys. *Appl Radiat Isotopes* 2011, **69**:1592–1595.

44. Jakubowski M, Palczynski C: **Beryllium in Handbook on the toxicology of metals.** Edited by Nordberg GF, Fowler BA, Nordberg M, Friberg LT. Amsterdam: Elsevier; 2007:415–431.
45. Fermo P, Beretta G, Maffei Facino R, Gelmini F, Piazzalunga A: **Ionic profile of honey as a potential indicator of botanical origin and global environmental pollution.** *Environ Pollut* 2013, **178**:173–181.
46. Association of Official Analytical Chemists (AOAC): *Official methods of analysis (15<sup>th</sup> edition).* Arlington: Association of Official Analytical Chemists Inc; 1990.
47. Pino A, Alimonti A, Botrè F, Minoia C, Bocca B, Conti ME: **Determination of twenty five elements in lichens by sector field inductively coupled plasma mass spectrometry and microwave assisted acid digestion.** *Rapid Commun Mass Sp* 2007, **21**:1900–1906.
48. Bocca B, Conti ME, Pino A, Mattei D, Forte G, Alimonti A: **Simple, fast, and low-contamination microwave-assisted digestion procedures for the determination of chemical elements in biological and environmental matrices by sector field ICP-MS.** *Int J Environ An Ch* 2007, **87**(15):1111–1123.
49. Conti ME, Muse JO, Mecozzi M: **Uncertainty in environmental analysis: theory and laboratory studies.** *Int J Risk Assess Manag* 2005, **5**:311–335.
50. Chudzinska M, Debska A, Baralkiewicz D: **Method validation for determination of 13 elements in honey samples by ICP-MS.** *Accred Qual Assur* 2012, **17**:65–73.
51. Royston P: **An extension of Shapiro and Wilk's W test for normality to large samples.** *Appl Stat* 1982, **31**:115–24.
52. Royston P: **Algorithm AS 181: The W test for Normality.** *Appl Stat* 1982, **31**:176–80.
53. Bauer DF: **Constructing confidence sets using rank statistics.** *J Am Stat Assoc* 1972, **67**:687–90.
54. Hollander M, Wolfe DA: *Nonparametric Statistical Methods.* New York: John Wiley & Sons; 1973:27–33. 68–75.
55. Conti ME, Iacobucci M, Cecchetti G: **A statistical approach applied to trace metal data from biomonitoring studies.** *Int J Environ Pollut* 2005, **23**(1):29–41.
56. Conti ME, Iacobucci M, Cucina D, Mecozzi M: **Multivariate statistical methods applied to biomonitoring studies.** *Int J Environ Pollut* 2007, **29**(1–3):333–343.
57. Szekely GJ, Rizzo ML: **Hierarchical Clustering via Joint Between-Within Distances: Extending Ward's Minimum Variance Method.** *J Classif* 2005, **22**(2):151–183.
58. Rizzo ML, Szekely GJ: **DISCO Analysis: A Nonparametric Extension of Analysis of Variance.** *Ann Appl Stat* 2010, **4**(2):1034–1055.
59. Szekely GJ, Rizzo ML: **Testing for Equal Distributions in High Dimension.** *Inter Stat* 2004, **5**.
60. Szekely GJ: *E-statistics: Energy of Statistical Samples, Department of Mathematics and Statistics, Bowling Green State University.* USA: Technical Report; 2000:03–05.
61. Thioulouse J, Chessel D, Doledec S, Olivier JM: **ADE-4: a multivariate analysis and graphical display software.** *Stat Comput* 1997, **7**:75–83.
62. Chessel D, Dufour A, Thioulouse J: **The ADE4 package. I. One-table methods.** *R News* 2004, **4**:5–10.
63. Conti ME, Mecozzi M: **Multivariate approaches to biomonitoring studies. Biological monitoring: Theory and applications. Bioindicators and biomarkers for environmental quality and human exposure assessment.** *The sustainable world M. E Conti (Ed)* 2008, **17**:213–226.
64. Conti ME, Finoia MG, Bocca B, Mele G, Alimonti A, Pino A: **Atmospheric background trace elements deposition in Tierra del Fuego region (Patagonia, Argentina), using transplanted *Usnea barbata* lichens.** *Environ Monit Assess* 2012, **184**:527–538.
65. Conti ME, Bocca B, Iacobucci M, Finoia MG, Mecozzi M, Pino A, Alimonti A: **Baseline trace metals in seagrass, algae and molluscs in a southern Tyrrhenian ecosystem (Linosa Island, Sicily).** *Arch Environ Con Tox* 2010, **58**:79–95.
66. Brereton RG: *Chemometrics: application of mathematics and statistics to laboratory systems.* Chichester, UK: Ellis Horwood; 1990.

doi:10.1186/1752-153X-8-44

**Cite this article as:** Conti et al.: Characterization of Argentine honeys on the basis of their mineral content and some typical quality parameters. *Chemistry Central Journal* 2014 **8**:44.

Publish with **ChemistryCentral** and every scientist can read your work free of charge

*“Open access provides opportunities to our colleagues in other parts of the globe, by allowing anyone to view the content free of charge.”*

W. Jeffery Hurst, The Hershey Company.

- available free of charge to the entire scientific community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
<http://www.chemistrycentral.com/manuscript/>

  
**ChemistryCentral**